DEPARTMENT OF OCEANOGRAPHY COLLEGE OF SCIENCES OLD DOMINION UNIVERSITY NORFOLK, VIRGINIA 23529

RESIDENCE TIME OF PARTICLE REACTIVE POLLUTANTS IN THE COASTAL SEA BED: CONTROL BY RESUSPENSION AND SEA BED MIXING PROCESSES

By

Donald J.P. Swift, Principal Investigator

Final Report For the period ended May 31, 1995

Prepared for Office of Naval Research Ballston Tower One 800 North Quincy St. Arlington, VA 22217-5660

Under

Research Contract N00014-93-1-0922 Joseph H. Kravitz, Scientific Officer Scientific Officer Code: 3242



Submitted by the Old Dominion University Research Foundation P.O. Box 6369
Norfolk, VA 23508-0369



March 1996

REPORT DOCUMENTATION PAGE

FORM APPROVED OMB No. 0704-0188

| Public reporting ourgen for this collection of gathering and maintaining the data needed of information, including suggestions for re Suite 1204, Arlington, VA 22202-4302 and | ducing the burden to Wash | ington Unada and S | Simulation: Send Comments regal | raing this buta | en estimate or any other asp | ita sources, sect of the collection Davis Highway, | |
|---|---|-------------------------------------|--|---------------------------------------|---|--|--|
| 1. AGENCY USE ONLY (Leave b | olank) 2. REPORT DATE 3. REPORT TYPE AND | | | | | | |
| 4. TITLE AND SUBTITLE OF REP POLLUTANTS IN THE COA AND SEA RED MIXING PR 6. AUTHOR(S) DONALD J.P. SWIFT, | STAL SEA BED: OCESSES: PRINCIPAL INV | CONTROL BY | ICLE REACTIVE RESUSPENSION | 1 | ING NUMBERS 14-93-1-0922 | ć | |
| 7. PERFORMING ORGANIZATION OLD DOMINION UNIVERSI P.O. BOX 6369 NORFOLK, VA 23508-030 DEPARTMENT OF OCEANOG | 59 RAPHY | | | 8. PERFO | RMING ORGANIZATI BER: | ON REPORT | |
| 9 SPONSORING MONITORING A OFFICE OF NAVAL RESEAT 800 NORTH QUINCY ST. ARLINGTON, VA 22217 | gency name(s) an RCH | ND ADDRESS(ES) | | 10. SPON REPO | SORING/MONITORIN RT NUMBER: | G AGENCY | |
| 11. SUPPLEMENTARY NOTES: | | | | | | | |
| UNCLASSIFIED; UNLIMITED | | | | 12b. DIST | 2b. DISTRIBUTION CODE | | |
| Our study was designed residence time of part methods for computing events, methods for codirectly from the spector establishing the rediffering half lives. | to develop made ticle-reactive the frequency omputing the details structure | pollutants and intensi iffusion coe | in the coastal s ty of resuspensi fficient for bio | eabed. on of bo logical | We have develottom sediment mixing of the | oped s by flow sea bed | |
| 14. SUBJECT TERMS | | | · · · · · · · · · · · · · · · · · · · | · · · · · · · · · · · · · · · · · · · | | | |
| - COSSEST TENNIS | | | | <u> </u> | 15. NUMBER OF PAC | 3ES: | |
| 7. SECURITY CLASSIFICATION OF REPORT: UNCLASSIFIED | EPORT: THIS PAGE | | 19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED | | 20. LIMITATION OF ABSTRACT UNCLASSIFIED | | |

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to stay within the lines to meet optical scanning requirements.

- Block 1. Agency Use Only (Leave blank).
- **Block 2.** Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.
- Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 30 Jun 88).
- Block 4. <u>Title and Subtitle</u>. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.
- Block 5. <u>Funding Numbers</u>. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract PR - Project G - Grant TA - Task

PE - Program WU - Work Unit Element Accession No.

- Block 6. <u>Author(s)</u>. Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).
- Block 7. <u>Performing Organization Name(s) and</u> Address(es). Self-explanatory.
- Block 8. <u>Performing Organization Report</u>
 <u>Number</u>. Enter the unique alphanumeric report
 number(s) assigned by the organization
 performing the report.
- Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.
- Block 10. Sponsoring/Monitoring Agency Report Number. (If known)
- Block 11. <u>Supplementary Notes</u>. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. <u>Distribution/Availability Statement</u>. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. Distribution Code.

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank. NTIS - Leave blank.

- **Block 13.** Abstract. Include a brief (Maximum 200 words) factual summary of the most significant information contained in the report.
- **Block 14.** <u>Subject Terms</u>. Keywords or phrases identifying major subjects in the report.
- **Block 15.** <u>Number of Pages</u>. Enter the total number of pages.
- **Block 16.** <u>Price Code</u>. Enter appropriate price code (NTIS only).
- Blocks 17. 19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.
- Block 20. <u>Limitation of Abstract</u>. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

RESIDENCE TIME OF PARTICLE REACTIVE POLLUTANTS IN THE COASTAL SEA BED: CONTROL BY RESUSPENSION AND SEA BED MIXING PROCESSES

Donald J. P. Swift, Alan W. Niedoroda (adjunct),* and G. T. F. Wong, Department of Oceanography, and D. Dauer, Department of Biology Old Dominion University

Norfolk, VA, 23504

Purpose of Study

Our study was designed to develop methods and algorithms that will allow us to predict the residence time of particle-reactive pollutants in the coastal seabed. We have developed methods for computing the frequency and intensity of resuspension of bottom sediments by flow events, methods for computing the diffusion coefficient for biological mixing of the sea bed directly from the species structure and population density of the benthic infauna, and methods for establishing the rate of sedimentation by comparison of profiles of radioisotope of differing half lives.

Background

Toxic pollutants entering estuarine and coastal waters are known to preferentially associate with the particulate phase, and their dispersal is therefore controlled by the natural cycle of fine sediment transport. A key aspect of this cycle is the exchange of sediment particles between the water column and the sea floor. As contaminated particles accumulate on the sea floor, they pass downward through a zone subject to biogenic mixing, and to resuspension by storm and tidal currents, with secondary release of the pollutant into the water column. Eventually, however, the sediment is so deeply buried that it is no longer accessible to these processes, and has have entered the zone of permanent burial. By measuring the rates of these competing processes (accumulation, mixing, resuspension), we will be able to predict hypothetical contaminant budgets under a variety of conditions.

Approach

In order to determine potential pattern of contaminant dispersal on the floor of southern Chesapeake Bay, we have undertaken fluid dynamical, radiogeochemical and biological investigations. We have developed a numerical model of boundary layer physics that will allows us to compute the frequency and intensity of particle resuspension, from wind, wave, and tidal current data. We have also collected cores in order to measure radionuclide profiles. The radionuclides ²³⁴Th, ¹³⁷Cs, and ²¹⁰Pb serve as proxies for the behavior of particle reactive pollutants. In addition, measurement of vertical concentration gradients of radionuclides with differing half lives has allowed us to estimate the other two critical rates, namely accumulation and mixing. We have also analyzed the composition of the benthic infaunal community as a function of depth. By applying random walk theory, we

^{*} Woodward Clyde Consultants 376 Hartfield Rd. Tallahassee FL 32303

have generated independent estimates of biodiffusive mixing. Finally, we have developed computational schemes that have allowed us to predict scenarios of contaminant release.

Results

Box cores and vibracores from the Baystem Plain of Southern Chesapeake Bay consist of a sandy clayey silt, containing a polychaete-molluscan community that extends, with vertically decreasing population density, to 30 cm depth. Calculations indicate depth-averaged biodiffusion coefficient for the upper 10 cm on the order of $1 \times 10^6 \, \mathrm{cm^2 y \, r^1}$. Radiographs indicate that the silt is well mixed in the summertime when winds are mild and the fauna is most active; in the Winter, however, the top several centimeters are frequently current stratified. Accumulation rates on the Baystem Plain range from 0 mm yr 1 near areas of tidal scour to 7 mm yr 1 in protected marginal bays. Computations indicate that the annual resuspension depth due to wave-tide interaction ranges from 0.4 to 1.6 cm in the shallower portions of the Baystem Plain (Fig. 1), and may be several times greater when wind-driven currents coincide with these interactions.

Burial histories for hypothetical contaminant layers vary from "rapid entombment" in marginal coves where accumulation is rapid relative to mixing, to "slow release" scenarios near the Bay Mouth, where both mixing and resuspension are intense. In a sample computation of such a scenario (Fig. 2) a site starts with an initial loading of the seabed of 0.8 mg of a hypothetical contaminant per cm² of the sea floor. As burial proceeds, much of the particle-reactive contamination leaks away through upward diffusion of the contaminant, coupled with repeated storm flushing of the surface layer. Eventually the contaminant layer is deeper than burrowers can reach, and the seabed contamination stabilizes at 5.6 mg of contaminant per cm of the sea floor.

Publications

- Niedoroda, A. W., , D. J. P. Swift, G. T. F. Wong, and D. Dauer, 1993 Residence time of particle-reactive pollutants in the estuary bed, Lower Chesapeake Bay: numerical simulation of resuspension and contaminant loss. Program with Abstracts, Estuarine Research Conference, Hilton Head, SC., November 16-18
- Swift, D. J. P., A. W. Niedoroda, G. T. F. Wong, and D. Dauer, 1993. Residence time of particle-reactive pollutants in the estuary bed, Lower Chesapeake Bay: Biology-based Estimates of sea bed mixing rates, Program with Abstracts, Estuarine Research Conference, Hilton Head, S.C.., November 16-18
- Niedoroda, A. W., Reed, C. and Swift, D. J. P., in preparation. Environmental management of contaminated sediments: factors controlling contamination, burial, and redistribution.




Fig. 1. ²¹⁰Pb inventories and accumulation rates in Southern Chesapeake Bay.




Fig. 2. This model run illustrates a burial history in which bioturbation is significant. For moderate to high values of the biodiffusion coefficient, surface concentration and surface flux approach low but appreciable limiting values with time. Total mass released increases continuously.